

HAMILTONIAN DIFFEOMORPHISMS OF TORIC MANIFOLDS

ANDRÉS VIÑA

ABSTRACT. We prove that $\pi_1(\text{Ham}(M))$ contains an infinite cyclic subgroup, where $\text{Ham}(M)$ is the Hamiltonian group of the one point blow up of CP^3 . We give a sufficient condition for the group $\pi_1(\text{Ham}(M))$ to contain an infinite cyclic subgroup, when M is a general toric manifold.

MSC 2000: 53D05, 57S05

1. INTRODUCTION

A loop ψ in the group $\text{Ham}(M, \omega)$ of Hamiltonian symplectomorphisms [9] of the symplectic manifold (M^{2n}, ω) determines a Hamiltonian fibration $E \xrightarrow{\pi} S^2$ with standard fibre M . On the total space E we can consider the first Chern class $c_1(VTE)$ of the vertical tangent bundle of E . Moreover on E is also defined the coupling class $c \in H^2(E, \mathbb{R})$ [6]. This class is determined by the following properties:

- i) $i_p^*(c)$ is the cohomology class of the symplectic structure on the fibre $\pi^{-1}(p)$, where i_p is the inclusion of $\pi^{-1}(p)$ in E and p is an arbitrary point of S^2 .
- ii) $c^{n+1} = 0$.

These canonical cohomology classes determine the characteristic number [7]

$$(1.1) \quad I_\psi = \int_E c_1(VTE)c^n.$$

I_ψ depends only on the homotopy class of ψ . Moreover I is an $\mathbb{R}$ -valued group homomorphism on $\pi_1(\text{Ham}(M, \omega))$, so the non vanishing of I implies that the group $\pi_1(\text{Ham}(M, \omega))$ is infinite. That is, I can be used to detect the infinitude of the corresponding homotopy group. Furthermore I calibrates the Hofer's norm ν on $\pi_1(\text{Ham}(M, \omega))$ in the sense that $\nu(\psi) \geq C|I_\psi|$, for all ψ , where C is a positive constant [12].

In [15] we gave an explicit expression for the value of the characteristic number I_ψ . This value can be calculated if one has a family of local symplectic trivializations of TM at one disposal, whose domains cover M and are fixed by the ψ_t 's. In fact we proved the following Theorem

Theorem 1. *Let $\psi : S^1 \rightarrow \text{Ham}(M, \omega)$ be a closed Hamiltonian isotopy. If $\{B_1, \dots, B_m\}$ is a set of symplectic trivializations for TM which covers M and such that $\psi_t(B_j) = B_j$, for all t and all j , then*

$$I_\psi = \sum_{i=1}^m J_i \int_{B_i \setminus \bigcup_{j < i} B_j} \omega^n + \sum_{i < k} N_{ik},$$

Key words and phrases. Hamiltonian Diffeomorphisms, toric manifolds, Symplectic Fibrations.

This work has been partially supported by Ministerio de Ciencia y Tecnología, grant MAT2003-09243-C02-00.

where

$$N_{ik} = n \frac{i}{2\pi} \int_{S^1} dt \int_{A_{ik}} (f_t \circ \psi_t)(d \log r_{ik}) \wedge \omega^{n-1},$$

$A_{ik} = (\partial B_i \setminus \cup_{r < k} B_r) \cap B_k$, J_i is the Maslov index of $(\psi_t)_*$ in the trivialization B_i and r_{ik} the corresponding transition function of $\det(TM)$.

The homotopy type of $\text{Ham}(M, \omega)$ is completely known in a few particular cases [8] [12] only. When M is a surface, $\text{Diff}_0(M)$ (the arc component of the identity map in the diffeomorphism group of M) is homotopy equivalent to the symplectomorphism group of M , so the topology of the groups $\text{Ham}(M)$ in dimension 2 can be deduced from the description of the diffeomorphism groups of surfaces given in [3] (see [12]). On the other hand, positivity of the intersections of J -holomorphic spheres in 4-manifolds have been used in [4] [1] [2] to prove results about the homotopy type of $\text{Ham}(M)$, when M is a ruled surface. But these arguments which work in dimension 2 or dimension 4 cannot be generalized to higher dimensions.

Let $\mathcal{O}$ be a coadjoint orbit of a Lie group G . If G is semisimple and acts effectively on $\mathcal{O}$, McDuff and Tolman have proved that the inclusion $G \rightarrow \text{Ham}(\mathcal{O})$ induces an injection from $\pi_1(G)$ to $\pi_1(\text{Ham}(\mathcal{O}))$ [11]. This result answers a question posed in [16]. In [14] we gave a lower bound for $\sharp\pi_1(\text{Ham}(\mathcal{O}))$, when $\mathcal{O}$ is a quantizable coadjoint orbit of a compact Lie group. In particular we proved that $\sharp\pi_1(\text{Ham}(\mathbb{C}P^n)) \geq n + 1$.

In this note we use Theorem 1 to prove that $\pi_1(\text{Ham}(M))$ contains an infinite cyclic subgroup, when M is a particular toric manifold. More precisely, when M is the 6-manifold associated to the polytope obtained truncating the tetrahedron of $\mathbb{R}^3$ with vertices $(0, 0, 0)$, $(\tau, 0, 0)$, $(0, \tau, 0)$, $(0, 0, \tau)$ by a horizontal plane [5]; that is, when M is the one point blow up of $\mathbb{C}P^3$. Moreover we will give a sufficient condition for $\pi_1(\text{Ham}(M))$ to contain an infinite cyclic subgroup, when M is a general toric manifold.

The paper is organized as follows. Section 2 is concerned with the determination of I_ψ for a natural circle action on the one point blow up of $\mathbb{C}P^3$. In Section 3 we generalize the arguments developed in Section 2 to toric manifolds. From this generalization it follows the aforesaid sufficient condition for the existence of an infinite subgroup in $\pi_1(\text{Ham}(M))$, when M is a toric manifold. Finally we check that this sufficient condition does not hold for $\mathbb{C}P^n$ with $n = 1, 2$. This is consistent with the fact that $\pi_1(\text{Ham}(\mathbb{C}P^n))$ is finite for $n = 1, 2$.

I thank Dusa McDuff for her enlightening comments.

2. HAMILTONIAN GROUP OF THE ONE POINT BLOW UP OF $\mathbb{C}P^3$

Given $\tau, \mu \in \mathbb{R}_{>0}$, with $\mu < \tau$, let M be the following manifold

$$(2.1) \quad M = \{z \in \mathbb{C}^5 : |z_1|^2 + |z_2|^2 + |z_3|^2 + |z_5|^2 = \tau/\pi, |z_3|^2 + |z_4|^2 = \mu/\pi\}/\mathbb{T}^2,$$

where the action of $\mathbb{T}^2$ is defined by

$$(2.2) \quad (a, b)(z_1, z_2, z_3, z_4, z_5) = (az_1, az_2, abz_3, bz_4, az_5),$$

for $a, b \in S^1$.

M is a toric 6-manifold; more precisely, it is the toric manifold associated to the polytope obtained truncating the tetrahedron of $\mathbb{R}^3$ with vertices

$$(0, 0, 0), (\tau, 0, 0), (0, \tau, 0), (0, 0, \tau)$$

by a horizontal plane through the point $(0, 0, \lambda)$, with $\lambda := \tau - \mu$ [5].

For $0 \neq z_j \in \mathbb{C}$ we put $z_j = \rho_j e^{i\theta_j}$, with $|z_j| = \rho_j$. On the set of points $[z] \in M$ with $z_i \neq 0$ for all i one can consider the coordinates

$$(2.3) \quad \left(\frac{\rho_1^2}{2}, \varphi_1, \frac{\rho_2^2}{2}, \varphi_2, \frac{\rho_3^2}{2}, \varphi_3 \right),$$

where the angle coordinates are defined by

$$(2.4) \quad \varphi_1 = \theta_1 - \theta_5, \varphi_2 = \theta_2 - \theta_5, \varphi_3 = \theta_3 - \theta_4 - \theta_5.$$

Then the standard symplectic structure on $\mathbb{C}^5$ induces the following form ω on this part of M

$$\omega = \sum_{j=1}^3 d\left(\frac{\rho_j^2}{2}\right) \wedge d\varphi_j.$$

Let $0 < \epsilon < 1$, we write

$$B_0 = \{[z] \in M : |z_j| > \epsilon, \text{ for all } j\}.$$

For a given $j \in \{1, 2, 3, 4, 5\}$ we set

$$B_j = \{[z] \in M : |z_j| < 2\epsilon \text{ and } |z_i| > \epsilon, \text{ for all } i \neq j\}$$

The family $B_0, \dots, B_5$ is not a covering of M , but if $[z] \notin \cup B_k$, then there are i, j , with $i \neq j$ and $|z_i| < \epsilon < |z_j|$.

On B_0 we will consider the well-defined Darboux coordinates (2.3). On B_1 , $\rho_j \neq 0$ for $j \neq 1$; so the angle coordinates φ_2 and φ_3 of (2.4) are well-defined. We put $x_1 + iy_1 := \rho_1 e^{i\varphi_1}$. In this way we take as symplectic coordinates on B_1

$$(x_1, y_1, \frac{\rho_2^2}{2}, \varphi_2, \frac{\rho_3^2}{2}, \varphi_3).$$

We will also consider the following Darboux coordinates: On B_2

$$\left(\frac{\rho_1^2}{2}, \varphi_1, x_2, y_2, \frac{\rho_3^2}{2}, \varphi_3 \right), \text{ with } x_2 + iy_2 := \rho_2 e^{i\varphi_2}.$$

On B_3

$$\left(\frac{\rho_1^2}{2}, \varphi_1, \frac{\rho_2^2}{2}, \varphi_2, x_3, y_3 \right), \text{ where } x_3 + iy_3 := \rho_3 e^{i\varphi_3}.$$

On B_4

$$\left(\frac{\rho_1^2}{2}, \varphi_1, \frac{\rho_2^2}{2}, \varphi_2, x_4, y_4 \right), \text{ with } x_4 + iy_4 := \rho_4 e^{i\varphi_4} \text{ and } \varphi_4 = \theta_4 - \theta_3 + \theta_5.$$

On B_5

$$(x_5, y_5, \frac{\rho_2^2}{2}, \chi_2, \frac{\rho_3^2}{2}, \chi_3),$$

where

$$x_5 + iy_5 := \rho_5 e^{i\varphi_5}, \chi_2 = \theta_2 - \theta_1, \chi_3 = \theta_3 - \theta_1 - \theta_4, \chi_5 = \theta_5 - \theta_1.$$

If $[z_1, \dots, z_5]$ is a point of

$$M \setminus \bigcup_{i=0}^5 B_i,$$

then there are $a \neq b \in \{1, \dots, 5\}$ such that $|z_a|, |z_b| < \epsilon$. We can cover the set $M \setminus \bigcup B_i$ by Darboux charts denoted $B_6, \dots, B_q$ similar to the preceding B_i 's

satisfying the following condition: The image of each B_a , with $a = 6, \dots, q$, is contained in a prism of $\mathbb{R}^6$ of the form

$$\prod_{i=1}^6 [c_i, d_i],$$

where at least four intervals $[c_i, d_i]$ have length of order ϵ .

By the infinitesimal “size” of the B_j , for $j \geq 1$, it turns out

$$(2.5) \quad \int_{B_j} \omega^3 = O(\epsilon), \text{ for } j \geq 1.$$

Let ψ_t be the symplectomorphism of M defined by

$$(2.6) \quad \psi_t[z] = [z_1 e^{2\pi i t}, z_2, z_3, z_4, z_5].$$

Then $\{\psi_t\}_t$ is a loop in the group $\text{Ham}(M)$ of Hamiltonian symplectomorphisms of M . By f is denoted the corresponding normalized Hamiltonian function. Hence $f = \pi \rho_1^2 - \kappa$ with $\kappa \in \mathbb{R}$ such that $\int_M f \omega^3 = 0$.

In the coordinates (2.3) of B_0 , ψ_t is the map $\varphi_1 \mapsto \varphi_1 + 2\pi t$. So the Maslov index $J_{B_0} = 0$. It follows from (2.5) and Theorem 1

$$(2.7) \quad I_\psi = \sum_{i < k} N_{ik} + O(\epsilon),$$

with

$$N_{ik} = \frac{3i}{2\pi} \int_{A_{ik}} f d \log r_{ik} \wedge \omega^2.$$

If $[z] \in A_{ik} \subset \partial B_i \cap B_k$, with $1 \leq i < k$, then at least the modules $|z_a|$ and $|z_b|$ of two components of $[z]$ are of order ϵ ; so N_{ik} is of order ϵ when $1 \leq i < k$. Analogously N_{0k} is of order ϵ , for $k = 6, \dots, q$. Hence (2.7) reduces to

$$(2.8) \quad I_\psi = \sum_{k=1}^5 N_{0k} + O(\epsilon).$$

If we put

$$(2.9) \quad N'_{0k} = \frac{3i}{2\pi} \int_{A'_{0k}} f d \log r_{ik} \wedge \omega^2,$$

with

$$A'_{0k} = \{[z] \in M : |z_k| = \epsilon, |z_r| > \epsilon \text{ for all } r \neq k\}$$

then

$$(2.10) \quad N_{0k} = N'_{0k} + O(\epsilon).$$

Next we determine the value of N'_{01} . To know the transition function r_{01} one needs the Jacobian matrix R of the transformation

$$(x_1, y_1, \frac{\rho_2^2}{2}, \varphi_2, \frac{\rho_3^2}{2}, \varphi_3) \rightarrow (\frac{\rho_1^2}{2}, \varphi_1, \frac{\rho_2^2}{2}, \varphi_2, \frac{\rho_3^2}{2}, \varphi_3)$$

in the points of A'_{01} ; where $\rho_1^2 = x_1^2 + y_1^2$, $\varphi_1 = \tan^{-1}(y_1/x_1)$. The function $r_{01} = \rho(R)$, where $\rho : Sp(6, \mathbb{R}) \rightarrow U(1)$ is the map which restricts to the determinant on $U(3)$ [13]. The non trivial block of R is the diagonal one

$$\begin{pmatrix} x_1 & y_1 \\ r & s \end{pmatrix},$$

with $r = -y_1(x_1^2 + y_1^2)^{-1}$ and $s = x_1(x_1^2 + y_1^2)^{-1}$. The non real eigenvalues of R are

$$\lambda_{\pm} = \frac{x_1 + s}{2} \pm \frac{i\sqrt{4 - (s + x_1)^2}}{2}.$$

On A'_{01} these non real eigenvalues occur when $(s + x_1)^2 < 2$, that is, if $|\cos \varphi_1| < 2\epsilon(\epsilon^2 + 1)^{-1} =: \delta$. If $y_1 > 0$ then λ_- of the first kind (see [13]) and λ_+ is of the first kind, if $y_1 < 0$.

Hence, on A'_{01} ,

$$\rho(R) = \begin{cases} \lambda_+ |\lambda_+|^{-1} = x + iy, & \text{if } |\cos \varphi_1| < \delta \text{ and } y_1 < 0; \\ \lambda_- |\lambda_-|^{-1} = x - iy, & \text{if } |\cos \varphi_1| < \delta \text{ and } y_1 > 0; \\ \pm 1, & \text{otherwise.} \end{cases}$$

where $x = \delta^{-1} \cos \varphi_1$, and $y = \sqrt{1 - x^2}$.

If we put $\rho(R) = e^{i\gamma}$, then $\cos \gamma = \delta^{-1} \cos \varphi_1$ (when $|\cos \varphi_1| < \delta$), and

$$\sin \gamma = \begin{cases} -\sqrt{1 - \cos^2 \gamma}, & \text{if } \sin \varphi_1 > 0; \\ \sqrt{1 - \cos^2 \gamma}, & \text{if } \sin \varphi_1 < 0. \end{cases}$$

So when φ_1 runs anticlockwise from 0 to 2π , γ goes round clockwise the circumference; that is, $\gamma = h(\varphi_1)$, where h is a function such that

$$(2.11) \quad h(0) = 2\pi, \text{ and } h(2\pi) = 0.$$

As $r_{01} = \rho(R)$, then $d \log r_{01} = idh$.

On A'_{01} the form ω reduces to $(1/2)d\rho_2^2 \wedge d\varphi_2 + d\rho_3^2 \wedge d\varphi_3$. From (2.9) one deduces

$$(2.12) \quad N'_{01} = \frac{3i}{4\pi} \int_{A'_{01}} if \frac{\partial h}{\partial \varphi_1} d\varphi_1 \wedge d\rho_2^2 \wedge d\varphi_2 \wedge d\rho_3^2 \wedge d\varphi_3.$$

The submanifold A'_{01} is oriented as a subset of ∂B_0 and the orientation of B_0 is the one defined by ω^3 , that is, by

$$d\rho_1^2 \wedge d\varphi_1 \wedge d\rho_2^2 \wedge d\varphi_2 \wedge d\rho_3^2 \wedge d\varphi_3.$$

Since $\rho_1 > \epsilon$ for the points of B_0 , then A'_{01} is oriented by $-d\varphi_1 \wedge d\rho_2^2 \wedge d\varphi_2 \wedge d\rho_3^2 \wedge d\varphi_3$. On the other hand, the Hamiltonian function $f = -\kappa + O(\epsilon)$ on A'_{01} . Then it follows from (2.12) together with (2.11)

$$N'_{01} = 6\pi^2 \kappa \int_0^{\mu/\pi} d\rho_3^2 \int_0^{\tau/\pi - \rho_3^2} d\rho_2^2 + O(\epsilon).$$

that is,

$$(2.13) \quad N'_{01} = 3\kappa(\tau^2 - \lambda^2) + O(\epsilon).$$

The contributions $N'_{02}, N'_{03}, N'_{04}, N'_{05}$ to I_ψ can be calculated in a similar way. One obtains the following results up to addends of order ϵ

$$(2.14) \quad N'_{02} = N_{05} = -(\tau^3 - \lambda^3) + 3\kappa(\tau^2 - \lambda^2), \quad N'_{03} = \tau^2(3\kappa - \tau), \quad N'_{04} = \lambda^2(3\kappa - \lambda).$$

As I_ψ is independent of ϵ , it follows from (2.8), (2.10), (2.13) and (2.14)

$$(2.15) \quad I_\psi = 6\kappa(2\tau^2 - \lambda^2) + \lambda^3 - 3\tau^3.$$

On the other hand, straightforward calculations give

$$\int_M \omega^3 = (\tau^3 - \lambda^3), \text{ and } \int_M \pi \rho_1^2 \omega^3 = \frac{1}{4}(\tau^4 - \lambda^4).$$

So

$$(2.16) \quad \kappa = \frac{1}{4} \left(\frac{\tau^4 - \lambda^4}{\tau^3 - \lambda^3} \right).$$

It follows from (2.15) and (2.16)

$$(2.17) \quad I_\psi = \frac{\lambda^2(-3\tau^4 + 8\tau^3\lambda - 6\tau^2\lambda^2 + \lambda^4)}{2(\tau^3 - \lambda^3)}.$$

Hence I_ψ is a rational function of τ and λ . It is easy to check that its numerator does not vanish for $0 < \lambda < \tau$. So we have

Proposition 2. *If ψ is the closed Hamiltonian isotopy defined in (2.6), then the characteristic number $I_\psi \neq 0$.*

Next we consider the loop $\tilde{\psi}$ defined by

$$(2.18) \quad \tilde{\psi}_t[z] = [z_1, z_2, z_3 e^{2\pi i t}, z_4, z_5].$$

The corresponding normalized Hamiltonian function is $\tilde{f} = \pi \rho_3^2 - \tilde{\kappa}$, where

$$(2.19) \quad \tilde{\kappa} = \frac{1}{4} \left(\frac{\tau^4 - 4\tau\lambda^3 + 3\lambda^4}{\tau^3 - \lambda^3} \right).$$

As in the preceding case

$$(2.20) \quad I_{\tilde{\psi}} = \sum_{j=1}^5 \tilde{N}'_{0j} + O(\epsilon),$$

where

$$\tilde{N}'_{0j} = \frac{3i}{2\pi} \int_{A'_{0j}} \tilde{f} d \log r_{0j} \wedge \omega^2.$$

The expression for $\tilde{N}'_{01}$ can be obtained from (2.12) substituting f for $\tilde{f}$; so

$$(2.21) \quad \tilde{N}'_{01} = -3(\tau - \tilde{\kappa})(\tau^2 - \lambda^2) + 2(\tau^3 - \lambda^3) + O(\epsilon).$$

Similar calculations give the following values for the $\tilde{N}'_{0j}$'s, up to summands of order ϵ ,

$$(2.22) \quad \tilde{N}'_{02} = \tilde{N}'_{05} = -3(\tau - \tilde{\kappa})(\tau^2 - \lambda^2) + 2(\tau^3 - \lambda^3), \quad \tilde{N}'_{03} = 3\tilde{\kappa}\tau^2, \quad \tilde{N}'_{04} = 3\lambda^2(\tilde{\kappa} - \mu).$$

It follows from (2.22), (2.21) and (2.20)

$$(2.23) \quad I_{\tilde{\psi}} = 6\tilde{\kappa}(2\tau^2 - \lambda^2) - 3(\tau^3 - 2\tau\lambda^2 + \lambda^3).$$

After (2.19) we obtain

$$I_{\tilde{\psi}} = -3I_\psi.$$

In the definition of M the variables z_1, z_2, z_5 play the same role. However we can consider the following S^1 action on M

$$(2.24) \quad \hat{\psi}_t[z] = [z_1, z_2, z_3, e^{2\pi i t} z_4, z_5].$$

Its Hamiltonian is $\hat{f} = \pi \rho_4^2 - \hat{\kappa}$, with

$$(2.25) \quad \hat{\kappa} = \frac{1}{4} \left(\frac{\lambda^4 - 4\lambda\tau^3 + 3\tau^4}{\tau^3 - \lambda^3} \right).$$

The corresponding $\hat{N}'_{0j}$ have the following values up to summand of order ϵ (2.26)
 $\hat{N}'_{01} = \hat{N}'_{02} = \hat{N}'_{05} = 3(\lambda + \hat{\kappa})(\tau^2 - \lambda^2) - 2(\tau^3 - \lambda^3)$, $\hat{N}'_{03} = 3\tau^2(\hat{\kappa} - \mu)$, $\hat{N}'_{04} = 3\hat{\kappa}\lambda^2$.
 From the preceding formulae one deduces

$$I_{\hat{\psi}} = -I_{\tilde{\psi}} = 3I_{\psi}.$$

Theorem 3. *Let M be the toric manifold defined by (2.1) and (2.2). If ψ , $\tilde{\psi}$ and $\hat{\psi}$ are the Hamiltonian loops in M defined by (2.6), (2.18) and (2.24) respectively, then*

$$I_{\hat{\psi}} = -I_{\tilde{\psi}} = 3I_{\psi},$$

with

$$I_{\psi} = \frac{\lambda^2(-3\tau^4 + 8\tau^3\lambda - 6\tau^2\lambda^2 + \lambda^4)}{2(\tau^3 - \lambda^3)},$$

λ being $\lambda := \tau - \mu$.

Corollary 4. *Let (M, ω) be the toric manifold one point blow up of $\mathbb{CP}^2$, then $\pi_1(\text{Ham}(M, \omega))$ contains an infinite cyclic subgroup.*

Proof. By Proposition 2 $I_{\psi} \neq 0$. As I is a group homomorphism then the class $[\psi^l] \in \pi_1(\text{Ham}(M, \omega))$ does not vanish, for all $l \in \mathbb{Z} \setminus \{0\}$. $\square$

3. HAMILTONIAN GROUP OF TORIC MANIFOLDS

In this Section we generalize the calculations carried out in Section 2 for the 6-manifold one point blow up of $\mathbb{CP}^3$ to a general toric manifold.

Let $\mathbb{T}$ be the torus $(S^1)^r$, and $\mathfrak{t} = \mathbb{R} \oplus \cdots \oplus \mathbb{R}$ its Lie algebra. Given $w_j \in \mathbb{Z}^r$, with $j = 1, \dots, m$ and $\tau \in \mathbb{R}^r$ we put

$$(3.1) \quad M = \{z \in \mathbb{C}^m : \pi \sum_{j=1}^m |z_j|^2 w_j = \tau\} / \mathbb{T},$$

where the relation defined by $\mathbb{T}$ is

$$(3.2) \quad (z_j) \simeq (z'_j) \text{ iff there is } \xi \in \mathfrak{t} \text{ such that } z'_j = z_j e^{2\pi i \langle w_j, \xi \rangle} \text{ for } j = 1, \dots, m.$$

We will assume that there is an open half space in $\mathbb{R}^r$ which contains all the vectors w_j and that $\{w_j\}_j$ span $\mathbb{R}^r$. We also assume that τ is a regular value of the map

$$z \in \mathbb{C}^m \mapsto \pi \sum_{j=1}^m |z_j|^2 w_j \in \mathbb{R}^r.$$

Then M is a closed toric manifold of dimension $n := 2(m - r)$ [10].

When $0 \neq z_a \in \mathbb{C}$, we write $z_a = \rho_a e^{i\theta_a}$. The standard symplectic form on $\mathbb{C}^m$ gives rise to the symplectic structure ω on M . On

$$\{[z] \in M : z_j \neq 0 \text{ for all } j\}$$

ω can be written as

$$\omega = \sum_{i=1}^n d\left(\frac{\rho_{ai}^2}{2}\right) \wedge d\varphi_{ai},$$

with φ_{ai} a linear combination of the θ_c 's.

Given $0 < \epsilon < 1$, we set

$$B_0 = \{[z] \in M : |z_j| > \epsilon \text{ for all } j\}$$

$$B_k = \{[z] \in M : |z_k| < \epsilon, |z_j| > \epsilon \text{ for all } j \neq k\},$$

as in Section 2. On B_0 we will consider the Darboux coordinates

$$\left\{ \frac{\rho_{ai}^2}{2}, \varphi_{ai} \right\}_{i=1, \dots, n}.$$

Given $k \in \{1, \dots, m\}$ we write ω in the form

$$\omega = d\left(\frac{\rho_k^2}{2}\right) \wedge d\varphi_k + \sum_{i=1}^{n-1} d\left(\frac{\rho_{ki}^2}{2}\right) \wedge d\varphi_{ki},$$

where φ_k and φ_{ki} are linear combinations of the θ_c 's. Then we consider on B_k the following Darboux coordinates

$$\left\{ x_k, y_k, \frac{\rho_{ki}^2}{2}, \varphi_{ki} \right\}_{i=1, \dots, n-1},$$

with $x_k + iy_k := \rho_k e^{i\varphi_k}$.

We denote by ψ_t the map

$$\psi_t : [z] \in M \mapsto [e^{2\pi it} z_1, z_2, \dots, z_m] \in M.$$

$\{\psi_t : t \in [0, 1]\}$ is a loop in $\text{Ham}(M)$. By repeating the arguments of Section 2 one obtains

$$I_\psi = \sum_{k=1}^m N'_{0k} + O(\epsilon),$$

where

$$N'_{0k} = \frac{ni}{2\pi} \int_{A'_{0k}} f d \log r_{0k} \wedge \omega^{n-1},$$

$$A'_{0k} = \{[z] \in M : |z_k| = \epsilon, |z_j| > \epsilon \text{ for all } j \neq k\},$$

and $f = \pi\rho_1^2 - \kappa$, with

$$\int_M \pi\rho_1^2 \omega^n = \kappa \int_M \omega^n.$$

As in Section 2, on A'_{0k} the exterior derivative $d \log r_{0k} = ih'(\varphi_k) d\varphi_k$, where $h = h(\varphi_k)$ is a function such that $h(0) = 2\pi$, $h(2\pi) = 0$. Then

$$N'_{0k} = -n \int_{\{[z] : z_k=0\}} f \omega^{n-1} + O(\epsilon).$$

Since I_ψ is independent of ϵ , we obtain

$$(3.3) \quad I_\psi = -n \sum_{k=1}^m \left(\int_{\{[z] : z_k=0\}} (\pi\rho_1^2 - \kappa) \omega^{n-1} \right).$$

This formula together with the fact that I is a group homomorphism give the following Theorem

Theorem 5. *Let (M, ω) be the toric manifold defined by (3.1) and (3.2). If*

$$\sum_{k=1}^m \left(\int_{\{[z] : z_k=0\}} (\pi\rho_1^2 - \kappa) \omega^{n-1} \right) \neq 0,$$

then $\pi_1(\text{Ham}(M, \omega))$ contains an infinite cyclic subgroup.

Examples. We will check the above result calculating I_ψ by (3.3) in two particular cases: When the manifold is $\mathbb{C}P^1$ and when is $\mathbb{C}P^2$.

For

$$\mathbb{C}P^1 = \{[z_1, z_2] : |z_1|^2 + |z_2|^2 = \tau/\pi\}/S^1$$

and $\psi_t[z_1, z_2] = [e^{2\pi it} z_1, z_2]$, the normalized Hamiltonian is $f = \pi\rho_1^2 - \tau/2$, that is, $\kappa = \tau/2$. In this case (3.3) reduces to $I_\psi = -\tau + 2\kappa = 0$. This is compatible with the fact that $\pi_1(\text{Ham}(\mathbb{C}P^1)) = \mathbb{Z}/2\mathbb{Z}$.

For

$$\mathbb{C}P^2 = \{[z_1, z_2, z_3] : |z_1|^2 + |z_2|^2 + |z_3|^2 = \tau/\pi\}/S^1,$$

the Hamiltonian is $f = \pi\rho_1^2 - \tau/3$. Moreover for $k \in \{1, 2, 3\}$

$$\int_{\{[z] : z_k=0\}} \omega = \tau.$$

On the other hand, for $k = 2, 3$

$$\int_{\{[z] : z_k=0\}} \pi\rho_1^2 \omega = \tau^2/2.$$

After (3.3) $I_\psi = -2(\tau^2 - 3\kappa\tau) = 0$. This result is consistent with the finiteness of $\pi_1(\text{Ham}(\mathbb{C}P^2))$, since $\text{Ham}(\mathbb{C}P^2)$ has the homotopy type of $PU(3)$ [4].

REFERENCES

- [1] M. Abreu, *Topology of symplectomorphism group of $S^2 \times S^2$* . Invent. Math. **131** (1998) 1-23
- [2] M. Abreu, D. McDuff, *Topology of symplectomorphism groups of rational ruled surfaces*. Journ. of Amer. Math. Soc. **13** (2000) 971-1009
- [3] C.J. Earle, J. Eells, *A fibre bundle description of Teichmüller theory*. J. Differential Geometry **3** (1969) 19-43
- [4] M. Gromov, *Pseudo holomorphic curves in symplectic manifolds*. Invent. Math. **82** (1985) 307-347
- [5] V. Guillemin, *Moment maps and combinatorial invariants of Hamiltonian T^n -spaces*. Birkhäuser, Boston, 1994
- [6] V. Guillemin, L. Lerman, S. Sternberg, *Symplectic fibrations and multiplicity diagrams*. Cambridge U.P., Cambridge, 1996
- [7] F. Lalonde, D. McDuff, L. Polterovich, *Topological rigidity of Hamiltonian loops and quantum homology*. Invent. Math. **135** (1999) 369-385
- [8] D. McDuff, *Lectures on groups of symplectomorphisms*. (Preprint [math.SG/0201032](#))
- [9] D. McDuff, D. Salamon, *Introduction to symplectic topology*, Clarendon Press, Oxford, 1998
- [10] D. McDuff, D. Salamon, *J-holomorphic curves and symplectic topology*, Amer. Math. Soc. Colloq. Publications, Providence, 2004
- [11] D. McDuff, S. Tolman, *On nearly semifree circle actions*. (Preprint [math.SG/0503467](#))
- [12] L. Polterovich, *The geometry of the group of symplectic diffeomorphisms*, Birkhäuser, Basel, 2001
- [13] D. Salamon, E. Zehnder, *Morse theory for periodic solutions of Hamiltonian systems and the Maslov index*. Comm. Pure Appl. Math **XLV**, 1303-1360 (1992)
- [14] A. Viña, *Generalized Symplectic Action and Symplectomorphism Groups of Coadjoint Orbits* (To appear in Ann. Global Anal. Geom.)
- [15] A. Viña, *A characteristic number of Hamiltonian bundles over S^2* . (Preprint [math.SG/0506174](#))
- [16] A. Weinstein, *Cohomology of symplectomorphism groups and critical values of Hamiltonians*. Math. Z. **201** (1989) 75-82

DEPARTAMENTO DE FÍSICA. UNIVERSIDAD DE OVIEDO. AVDA CALVO SOTELO. 33007 OVIEDO. SPAIN.

E-mail address: vina@uniovi.es